



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

191. Proposed by J. EDWARD SANDERS, Reinersville, Ohio.

Two random lines cut a given circle. What is the chance that they intersect within the circle?

Solution by HENRY HEATON, Belfield, N. D., and the PROPOSER.

Let x = the distance of one of the lines from the center of the circle, and θ = the angle between the lines. The length of the part of the first line lying within the circle is $2\sqrt{(a^2 - x^2)}$. For given values of x and θ the chance of intersection is

$$\frac{2\sin \theta \sqrt{(a^2 - x^2)}}{2a} = \frac{\sin \theta}{a} \sqrt{(a^2 - x^2)},$$

and the required chance is

$$P = \int_0^a \int_0^{4\pi} \frac{\sin \theta}{a} \sqrt{(a^2 - x^2)} dx d\theta \div \int_0^a \int_0^{4\pi} dx d\theta = \frac{2}{a\pi} \int_0^a \sqrt{(a^2 - x^2)} dx = \frac{1}{2}.$$

Also solved by G. B. M. Zerr, who gets $\frac{1}{2}$ for the result. His solution will be published in the next issue of the MONTHLY.

PROBLEMS FOR SOLUTION.

ALGEBRA.

293. Proposed by C. E. WHITE, Vanderbilt University, Nashville, Tenn.

Prove by mathematical induction that $\frac{(x-a)^{m-1}}{(m-1)!} f^{m-1}(a) + \frac{(x-a)^{m-2}}{(m-2)!} f^{m-2}(a) + \dots + \frac{(x-a)^2}{2!} f''(a) + (x-a)f'(a) + f(a)$ will be the remainder when $f(x)$ is divided by $(x-a)^m$.

294. Proposed by O. L. CALLECOT, Gettysburg, S. Dak.

Find the limit of $\sum_{n=1}^{\infty} \frac{2(n^2 + 3n + 3)}{n(n+1)(n+2)(n+3)}$.

GEOMETRY.

326. Proposed by L. E. NEWCOMB, Los Gatos, Calif.

The circle C of radius pR encloses the circles A_1B_1 of radii R and $(p-1)R$, respectively; the circle B_1 is tangent to $A_1B_1C_1$; the circle B_2 is tangent to AB_1C ; the circle B_3 to AB_2C , ..., B_n to $AB_{n-1}C$. Find the radius of the circle B_n .

327. Proposed by J. C. CORBIN, Pine Bluff, Ark.

In triangle ABC , the triangle DEF is formed by joining the feet of the medians and four parallelograms are also formed, viz., $AEDF$, $BFED$, and $CEFD$. Let a, b, c, d, e, f represent the three medians of ABC , and the three sides of DEF . Then the sum of the squares of the six diagonals equals the sum of the squares of the twelve sides of the parallelograms, which are equal in sets of four. That is, $a^2 + b^2 + c^2 + d^2 + e^2 + f^2 = 4(d^2 + e^2 + f^2)$, or $a^2 + b^2 + c^2 = 3(d^2 + e^2 + f^2) = 3/4(AB^2 + BC^2 + CA^2)$.

328. Proposed by CHARLES GILPIN, JR., Philadelphia, Pa.

A sphere with the radius R is divided into two segments by a plane passed through it half way between the center and circumference. The smaller segment is divided into two parts by a plane passed through it at right angles to the base and cutting it half way between its center and circumference. Required the contents of the two parts of the segment.

CALCULUS.

250. Proposed by V. M. SPUNAR, East Pittsburg, Pa.

Differentiate $(\log^n x)$.

251. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Find in terms of x the functions c_1x and c_2x defined, respectively, by the relations

$$\begin{aligned} (a) \quad x \log(c_1x) &= c_1x \log x, \\ (b) \quad x \log x &= c_2x \log(c_2x). \end{aligned}$$

MECHANICS.

210. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

A rigid triangle is formed of three weightless, smoothly jointed, rigid rods BC, CA, AB . At their mid points D, E, F , respectively, are small, smooth rings, through which passes an endless, stretched, elastic string, forming the triangle DEF . Find by graphical construction the reaction at the joints.

211. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

A smooth elliptic wire, axis vertical, has a small ring sliding on it, connected by elastic strings with each focus. Either string is just unstretched when the ring is nearest the corresponding focus. The modulus of elasticity is W/n , where W oz. is the weight of the ring. Find the distance of the ring from the upper focus in the different positions of equilibrium, and in each case discuss the nature of the equilibrium.

MISCELLANEOUS.

175. Proposed by PROFESSOR R. D. CARMICHAEL, Anniston, Ala.

If x and z are connected by the relation $z = zf(x) + x\phi(z)$, find the value of $f(z)$ in the form of a power series in x with constant coefficients. In particular, give such a value of z when $z = z \sin x + x \cos z$.